


Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Southwest Florida Monday, 15 June 2015 NOAA National Ocean Service NOAA Satellite and Information Service NOAA National Weather Service Last bulletin: Monday, June 8, 2015



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from June 5 to 10: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Florida Fish and Wildlife Conservation Commission (FWC) Fish and Wildlife Research Institute. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through FWC Fish and Wildlife Research Institute at: http://myfwc.com/redtidestatus

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit at: $\frac{\text{http://tidesandcurrents.noaa.gov/hab/bulletins.html}}{\text{http://tidesandcurrents.noaa.gov/hab/bulletins.html}}$

Conditions Report

There is currently no indication of *Karenia brevis* (commonly known as Florida red tide) along the coast of southwest Florida, including the Florida Keys. No respiratory irritation is expected alongshore southwest Florida Monday, June 15 through Monday, June 22. Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

Recent samples received from along- and offshore southwest Florida, from Pinellas to Monroe counties, including the Florida Keys, all indicate that *Karenia brevis* is not present (FWRI, SCHD, MML, CCPCPD; 6/8-6/11).

Recent MODIS Aqua imagery (6/13, shown left) is completely obscured by clouds alongand offshore southwest Florida from Lee County to the Florida Keys, preventing analysis. Elevated chlorophyll (1-5 μ g/L) is visible in patches along- and offshore southwest Florida from Pinellas to Charlotte counties. The elevated chlorophyll is likely associated with blooms of various algal species that continue to be detected alongshore southwest Florida.

Harmful algal bloom formation at the coast of southwest Florida is not expected today through Monday, June 22.

Davis, Yang




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

-2-

Wind Analysis

Englewood to Tarpon Springs (Venice): Southeast winds (5-15kn, 3-8m/s) today. West winds (5kn, 3m/s) tonight becoming east winds (10kn, 5m/s) after midnight. East winds (10kn) Tuesday becoming southeast winds (5kn) in the afternoon. North to northeast winds (10kn) Tuesday night. East to southeast winds (5kn) Wednesday. Southwest winds (5kn) Wednesday night becoming southeast winds (10kn) after midnight. Southeast to south winds (10kn) Thursday becoming southwest winds (5kn) Thursday night. Southeast to south winds (10kn) Friday.



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).